

221725  
 221726  
 221727  
 221728  
 221729  
 221730  
 221731  
 221732  
 221733  
 221734  
 221735  
 221736  
 221737  
 221738  
 221739  
 221740  
 221741  
 221742  
 221743  
 221744  
 221745  
 221746  
 221747  
 221748  
 221749  
 221750  
 221751  
 221752  
 221753  
 221754  
 221755  
 221756  
 221757  
 221758  
 221759  
 221760  
 221761  
 221762  
 221763  
 221764  
 221765  
 221766  
 221767  
 221768  
 221769  
 221770  
 221771  
 221772  
 221773  
 221774  
 221775  
 221776  
 221777  
 221778  
 221779  
 221780  
 221781  
 221782  
 221783  
 221784  
 221785  
 221786  
 221787  
 221788  
 221789  
 221790  
 221791  
 221792  
 221793  
 221794  
 221795  
 221796  
 221797  
 221798  
 221799  
 221800

2. A method according to claim 1 (b) wherein the known ligand-responsive receptor protein is selected from the group consisting of glucocorticoid receptor, mineralocorticoid receptor, human thyroid receptors alpha and beta and rat thyroid receptor alpha, estrogen-related receptors hERR1 and hERR2, and retinoic acid receptors alpha and beta.

4. A method according to claim 1 (c) (2) wherein the reporter gene is selected from the group consisting of a chloramphenicol acetyltransferase (CAT) gene and a firefly luciferase gene.

6. A method according to claim 5 wherein the glucocorticoid response element is encompassed within the mammary tumor virus long terminal repeat sequence (MTV LTR), and the thyroid response element is encompassed within the growth hormone promoter sequence.

(a) an expressible chimeric DNA sequence (c) comprised of operative portions of a DNA-binding domain of a first receptor sequence linked to operative portions of a ligand-binding domain of a second receptor sequence, and (b) a reporter nucleic acid sequence functionally linked to an operative hormone response element wherein said chimeric DNA sequence is expressed and wherein the DNA-binding domain of the chimeric receptor protein thus produced can functionally bind to and activate the hormone response element that is functionally linked to the reporter sequence.

28

9. A method according to claim 7 wherein the reporter gene is selected from the group consisting of a chloramphenicol acetyltransferase (CAT) gene and a firefly luciferase gene.

**11.** A method according to claim 10 wherein the glucocorticoid response element is encompassed within the mammary tumor virus long terminal repeat sequence (MTV LTR), and the thyroid response element is encompassed within the growth hormone promoter sequence.

\* \* \* \* \*

12. A method for identifying functional ligands for receptor proteins, said method comprising:

(a) isolating DNA sequences having a ligand-binding domain and a DNA-binding domain;

(b) constructing a chimeric gene by substituting operative portions of the DNA-binding domain region of the DNA sequence of step (a) with operative portions of a DNA-binding domain region from a known ligand-responsive receptor protein selected from the group consisting of hGR, hMR, hERR, hERR<sub>1</sub>, rTR $\alpha$ , hT<sub>1</sub>R $\alpha$ , hT<sub>1</sub>R $\beta$ , hRAR $\alpha$ 1 and hRAR $\beta$ ;

(c) introducing into a suitable receptor-deficient host cell: (1) the chimeric gene from step (b), and (2) a reporter gene functionally linked to an operative hormone response element wherein the hormone response element is capable of being activated by the DNA-binding domain region of the receptor protein encoded by the chimeric gene of step (b) and is selected from the group consisting of wild-type, recombinantly produced or synthetic GR, PR, ER, T<sub>1</sub>R $\beta$  and V-erbA response element;

(d) challenging the transfected host cell from step (c) with at least one compound to be evaluated for ligand binding activity with the chimeric receptor protein encoded by the chimeric gene of step (b);

(e) monitoring induction of the reporter gene; and

(f) identifying as a functional ligand(s) that ligand(s) which induces production of the protein product of the reporter gene.

13. A method according to claim 12 wherein the glucocorticoid response element is encompassed within the mammary tumor virus long terminal repeat sequence (MTV LTR), and the thyroid response element is encompassed within the growth hormone promoter sequence.

14. A method for identifying functional ligands for receptor proteins in a cell wherein said cell contains: (a) an expressible chimeric DNA sequence (c) comprised of operative portions of a DNA-binding domain of a first receptor sequence linked to operative portions of a ligand-binding domain of a second receptor sequence, wherein said first and second receptor sequences are selected from the group consisting of sequences from hGR, hMR, hERR, hERR<sub>1</sub>, rTR $\alpha$ , hT<sub>1</sub>R $\alpha$ , hT<sub>1</sub>R $\beta$ , hRAR $\alpha$ 1 and hRAR $\beta$ , and (b) a reporter nucleic acid sequence functionally linked to an operative hormone response element wherein said chimeric DNA sequence is expressed and wherein the DNA-binding domain of the chimeric receptor protein thus produced can functionally bind to and activate the hormone response element that is functionally linked to the reporter sequence, wherein the hormone response element is selected from the group consisting of wild-type, recombinantly produced or synthetic GR, PR, ER, T<sub>1</sub>R $\beta$  and V-erbA response element; said method comprising challenging the cell with at least one compound to be evaluated for ligand binding activity wherein said compound to be evaluated is not known to be a functional ligand for the chimeric protein encoded by said chimeric DNA sequence (c).

15. A method according to claim 14 wherein the glucocorticoid response element is encompassed within the mammary tumor virus long terminal repeat sequence (MTV LTR), and the thyroid response element is encompassed within the growth hormone promoter sequence.

FOIA b7 - D